

## ***Efficient Elliptic Curve Cryptosystems Using Efficient Exponentiation***

Kamal Darweesh, Mohammad Saleh\*

*Mathematics Department & Scientific Computing Master Program, Birzeit University,  
Palestine*

---

### **Abstract**

Elliptic curve cryptosystem (ECC) is a new generation of public key cryptosystems that has smaller key size for the same level of security. The exponentiation in elliptic curve is the most important operation in ECC, so when the ECC is put into practice, the major problem is how to enhance the speed of the exponentiation. It is thus of great interest to develop algorithms for exponentiation, which allow efficient implementations of ECC.

In this paper, we improve efficient algorithm for exponentiation on elliptic curve defined over  $\mathbf{F}_p$  in terms of affine coordinates. The algorithm computes  $2^{n_2} (2^{n_1} P + Q)$  directly from random points P and Q on an elliptic curve, without computing the intermediate points. Moreover, we apply the algorithm to exponentiation on elliptic curve with Width-w Mutual Opposite Form (wMOF) and analyze their computational complexity. This algorithm can speed the wMOF exponentiation of elliptic curve of size 160-bit about (21.7%) as a result of its implementation with respect to affine coordinates.

---

### **1. Introduction**

Elliptic curve cryptosystem, which were suggested independently by Miller [7] and Koblitz, it is a new generation of public key cryptosystems that has smaller key sizes for the same level of security.

The elliptic curve cryptographic operations, like encryption/decryption schemes generation/verification signature, require computing of exponentiation on elliptic curve. The computational performance of elliptic curve cryptographic protocol such as Diffie-Hellman [4] Key Exchange protocol strongly depends on the efficiency of exponentiation, because it is the costliest operation. Therefore, it is very attractive to speed up exponentiation by providing algorithms that allow efficient implementations of elliptic curve cryptosystems [2, 5, 6, 8, 9, 12].

Binary method is one of typical methods that can speed up exponentiation by reducing additions, where addition of two points and doubling of two points are performed repeatedly. It is based on the binary representation of the exponent, so the

---

\* Corresponding author, Email: [msaleh@birzeit.edu](mailto:msaleh@birzeit.edu)

Copyright © 2007 IeJs-IeJEMTA. <http://www.iejs.org/>

average number of addition of elliptic points operations required by the binary method depends on the minimal hamming weight of the exponent.

wMOF is a base-2 representation which provide the minimal hamming weight of exponent. Their great advantage is that they can be generated from left-to-right which means, that the recoding doesn't have to be done in a separate stage, but can be performed on-the-fly during the evaluation. As a result, it is no longer necessary to store the whole recoded exponent, but only small parts at once.

Another approach to speed up exponentiation is by increasing the speed of doublings. One method to speed the doublings is direct computation of several doubling, which computes  $2^n P$  directly from  $P \in E(\mathbf{F}_q)$ , without computing intermediate points  $2P, 2^2P, \dots, 2^{n-1}P$ . Sakai and Sakurai [12] proposed formulae for computing  $2^n P$  directly ( $\forall n \geq 1$ ) on  $E(\mathbf{F}_p)$  in terms of affine coordinates. Since modular inversion is more expensive than multiplication, their formula requires only one inversion for computing  $2^n P$  instead of  $n$  inversions in usual add-double method.

In this paper, we construct efficient formula to compute  $2^{n_2} (2^{n_1} P + Q)$  directly from  $P, Q \in E(\mathbf{F}_p)$ , without computing intermediate points  $2P, 2^2P, \dots, 2^{n_1} P$ ,  $2(2^{n_1} P + Q), \dots, 2^{n_2-1}(2^{n_1} P + Q)$ , where  $n_j \geq 1$ . Our formula has computational complexity  $(4n+10)M+(4n+6)S+I$ , where  $M$ ,  $S$  and  $I$  denote multiplication, squaring and inversion respectively in  $\mathbf{F}_p$ , and  $n=n_1+n_2$ .

Moreover, we show in which way this new algorithm for direct computing  $2^{n_2} (2^{n_1} P + Q)$  can be combined with wMOF exponentiation method [11]. We also implement wMOF exponentiation with and without this formula and discuss the efficiency. The result of this implementation shows that 21.7% speed increase in wMOF exponentiation with this formula on elliptic curve of size 160-bit.

Let  $\mathbf{F}_p$  denotes a prime finite field with  $p$  elements.

We consider an elliptic curve  $E$  given by Weierstrass non-homogeneous equation:

$$E: y^2 = x^3 + ax + b$$

Where  $a, b \in \mathbf{F}_p$ ,  $p > 3$ , and  $4a^3 + 27b^2 \neq 0$  (i.e.  $E$  is smooth).

Let  $P_1 = (x_1, y_1)$ ,  $P'_1 = (x'_1, y'_1)$ ,  $P_{2^n} = 2^n P_1 = (x_{2^n}, y_{2^n}) \in E(\mathbf{F}_p)$ .

Let the elliptic curve point addition and doubling be denoted by ECADD and ECDBL, respectively. Let  $M$ ,  $S$  and  $I$  denote multiplication, squaring and inversion, respectively in  $\mathbf{F}_p$ , where  $S=0.8M$ , as it is customary nowadays. Therefore, inversions are very costly compared to multiplications and squaring and should be avoided.

This paper is organized as follows: In Sect. 1, we give some definitions and notations. In Sect. 2, we summarize pervious work. In Sect. 3, we will describe our algorithm for direct computing of  $2^{n_2} (2^{n_1} P + Q)$  in terms of affine coordinates. In Sect. 4, we use this formula in exponentiation with wMOF method, and show in what way this new derived formula can improve the speed of the exponentiation. In Sect. 5 timing of our implementation will be given. Finally conclusions will be given in Sect. 6.

## 2. Previous work

In this section, we summarize the known algorithms for point addition, point doublings, and direct doublings.

### 2.1 Point addition

In terms of affine coordinates, point addition can be computed as follows:

Let  $P_1 = (x_1, y_1)$ , and  $Q = (x, y)$ ,  $\neq O$  where  $O$  denotes the point at infinity, then  $P' = (x', y')$  can be computed as follows

$$\begin{aligned} x' &= \lambda^2 - x_1 - x \\ y' &= \lambda(x_1 - x') - y_1 \\ \lambda &= \frac{(y - y_1)}{(x - x_1)} \end{aligned} \quad (2.1)$$

The formulae above have computational complexity  $S + 2M + I$  [1]

### 2.2 Point doubling

In terms of affine coordinates, point addition can be computed as follows:

Assume Let  $P_1 = (x_1, y_1) \neq O$  where  $O$  denotes the point at infinity, then  $2P = P_2 = (x_2, y_2)$  can be computed as follows

$$\begin{aligned} x_2 &= \lambda^2 - 2x_1 \\ y_2 &= \lambda(x_1 - x_2) - y_1 \\ \lambda &= \frac{3x_1^2 + a}{2y_1} \end{aligned} \quad (2.2)$$

The formulae above have computational complexity  $2S + 2M + I$  [2]

### 2.3 Direct Doubling

One method to increase the speed of doublings is direct computation of several doublings, which can compute  $2^n P$  directly from  $P \in E(\mathbf{F}_q)$ , without computing the intermediate points  $2P, 2^2P, \dots, 2^{n-1}P$  [12].

Guajardo and Paar [5] suggested increase doubling speed by formulating algorithms for direct computation of  $4P$ ,  $8P$ , and  $16P$  on elliptic curves over  $\mathbf{F}_2^m$  in terms of affine coordinates. Sakai and Sakurai [12] proposed formulae for computing  $2^n P$  directly ( $\forall n \geq 1$ ) on  $E(\mathbf{F}_p)$  in terms of affine coordinates.

These formulas require only one inversion for computing  $2^n P$  instead of  $d$  inversions in regular add-double method. Therefore direct computation of several doublings may be effective in elliptic curve exponentiation because modular inversion is more expensive than multiplication.

## 3. Direct Computation of $2^{n_2} (2^{n_1} P + Q)$ in affine coordinate

In this paper, we derive formula for computing  $2^{n_2} (2^{n_1} P + Q)$  directly from a given point  $P, Q \in E(\mathbf{F}_p)$  without computing the intermediate points  $2P, 2^2P, \dots, 2^{n_1} P$ ,

$2(2^{n_1}P+Q), \dots, 2^{n_2-1}(2^{n_1}P+Q)$ , where  $n_i \geq 1$ , in terms of affine coordinate. This formula can work with wMOF exponentiation method.

We begin by constructing formula for small  $n_1, n_2$ , then we will construct algorithm for general  $n_1, n_2$ .

As an example, let  $n_1 = 2, n_2 = 1$ , let  $P_1 = (x_1, y_1), Q = (x, y), P'_1 = (x'_1, y'_1) \in E(\mathbb{F}_p)$  then for an elliptic curve with weierstrass form in terms of affine coordinates  $P'_2 = 2P'_1 = 2(4P_1+Q) = (x'_2, y'_2)$  can be computed as the following:

(1) Computing  $4P_1$  as in [12]

$4P_1 = P_4 = (x_4, y_4)$  can be computed as follows:

$$x_4 = \frac{A_2}{(4C_0C_1)^2} \quad (3.1)$$

$$y_4 = \frac{C_2}{(4C_0C_1)^3} \quad (3.2)$$

(2) Computing  $(4P_1+Q)$

Assume  $4P_1 = (x_4, y_4) \neq -Q$ , recall from Sect. 2.1, the point addition then  $P'_1 = (x'_1, y'_1) = (4P_1+Q)$  in term of affine coordinates, can be computed as follows:

$$\lambda = \frac{C_2 - (4C_0C_1)^3 y}{(4C_0C_1)(A_2 - (4C_0C_1)^2 x)} \quad (3.3)$$

Now let

$T = C_2 - (4C_0C_1)^3 y, S = A_2 - (4C_0C_1)^2 x$ , we get:

$$\lambda = \frac{T}{(4C_0C_1)S} \quad (3.4)$$

Substituting  $\lambda$ , and  $x_4$  into the expression for  $x'_1$ , we find

$$x'_1 = \frac{T^2}{(4C_0C_1)^2 S^2} - x - \frac{A_2}{(4C_0C_1)^2} \quad (3.5)$$

After simplification equation (3.5) we get:

$$x'_1 = \frac{T^2 - S^2(A_2 + (4C_0C_1)^2 x)}{(4C_0C_1)^2 S^2} \quad (3.6)$$

Let  $M = A_2 + (4C_0C_1)^2 x$ , we get :

$$x'_1 = \frac{T^2 - MS^2}{(4C_0C_1)^2 S^2} \quad (3.7)$$

Let  $A'_0 = T^2 - MS^2$ , we get:

$$x'_1 = \frac{A'_0}{(4C_0C_1)^2S^2} \quad (3.8)$$

Substituting  $\lambda$ , and  $x'_1$  into the expression for  $y'_1$ , we find

$$y'_1 = \frac{T}{(4C_0C_1)S} \left( x - \frac{A'_0}{(4C_0C_1)^2S^2} \right) - y \quad (3.9)$$

After simplification we get:

$$y'_1 = \frac{-(4C_0C_1)^3yS^3 - T(A'_0 - (4C_0C_1)^2xS^2)}{(4C_0C_1)^3S^3} \quad (3.10)$$

Let  $C'_0 = -(4C_0C_1)^3yS^3 - T(A'_0 - (4C_0C_1)^2xS^2)$ , we get:

$$y'_1 = \frac{C'_0}{(4C_0C_1)^3S^3} \quad (3.11)$$

(3) Computing  $2(4P_1+Q)=2P'_1$

Recall from Sect. 2.2, the point doubling, then  $2P'_1 = P'_2 = (x'_2, y'_2)$  in term of affine coordinates, can be computed as follows:

$$\lambda = \frac{3 \left( \frac{A'_0}{(4C_0C_1)^2S^2} \right)^2 + a}{2 \left( \frac{C'_0}{(4C_0C_1)^3S^3} \right)} \quad (3.12)$$

After simplification we get:

$$\lambda = \frac{3A'^2_0 + a(4C_0C_1)^4S^4}{2C'_0(4C_0C_1)S} \quad (3.13)$$

Now, let  $B'_0 = 3A'^2_0 + a(4C_0C_1)^4S^4$ , we get:

$$\lambda = \frac{B'_0}{2C'_0(4C_0C_1)S} \quad (3.14)$$

Substituting  $\lambda$ , and  $x'_1$  into the expression for  $x'_2$ , we find

$$x'_2 = \frac{B'^2_0}{(2C'_0)^2(4C_0C_1)^2S^2} - 2 \left( \frac{A'_0}{(4C_0C_1)^2S^2} \right) \quad (3.15)$$

After simplification we get:

$$x'_2 = \frac{B'^2_0 - 8A'_0C'^2_0}{(2C'_0)^2(4C_0C_1)^2S^2} \quad (3.16)$$

Let  $A'_1 = B'^2_0 - 8A'_0C'^2_0$ , we get:

$$x'_2 = \frac{A'_1}{(2C'_0)^2(4C_0C_1)^2S^2} \quad (3.17)$$

Substituting  $\lambda$ ,  $y'_1$ ,  $x'_1$  and  $x'_2$  into the expression for  $y'_2$ , we find

$$y'_2 = \frac{B'_0}{2C'_0(4C_0C_1)S} \left( \frac{A'_0}{(4C_0C_1)^2S^2} \right) - \left( \frac{A'_1}{(2C'_0)^2(4C_0C_1)^2S^2} \right) - \frac{C'_0}{(4C_0C_1)^3S^3} \quad (3.18)$$

After simplification we get:

$$y'_2 = \frac{-8C'_0{}^4 - B'_0(A'_1 - 4A'_0C'_0{}^2)}{(2C'_0)^3(4C_0C_1)^3S^3} \quad (3.19)$$

Let  $C'_1 = -8C'_0{}^4 - B'_0(A'_1 - 4A'_0C'_0{}^2)$ , we get finally:

$$y'_2 = \frac{C'_1}{(2C'_0)^3(4C_0C_1)^3S^3} \quad (3.20)$$

The formulae above have computational complexity  $18S + 22M + I$

### 3.1 The formulae Computing $2^{n_2}(2^{n_1}P + Q)$ in Affine Coordinate

From above formulae for direct computing  $2(4P_1+Q)$ , we can easily obtain general formulae that allow direct computing  $2^{n_2}(2^{n_1}P + Q)$  for  $n_1 \geq 1$ . Algorithm 3.1 describes these formulae.

---

**Algorithm 3.1 Direct Computation of  $2^{n_2}(2^{n_1}P + Q)$  in affine coordinate, where  $n_1 \geq 1$ , and  $P, Q \in E(\mathbf{F}_p)$ .**

---

INPUT:  $P_1 = (x_1, y_1), Q = (x, y) \in E(\mathbf{F}_p)$

OUTPUT:  $P'_{2^4} = 2^4P' = 2^4(2P_1+Q) = (x'_{2^4}, y'_{2^4}) \in E(\mathbf{F}_p)$

1. Compute  $A_0$  and  $C_0$  and  $B_0$

$$C_0 = y_1$$

$$A_0 = x_1$$

$$B_0 = 3x_1^2 + a$$

2. For  $i$  from 1 to  $n_1$  Compute  $A_i, C_i$ , for  $i$  from 1 to  $n_1 - 1$  Compute  $B_i$

$$A_i = B_{i-1}^2 - 8A_{i-1}C_{i-1}^2$$

$$C_i = -8C_{i-1}^4 - B_{i-1}(A_i - 4A_{i-1}C_{i-1}^2)$$

$$B_i = 3A_i^2 + 16^i a \left( \prod_{j=0}^{i-1} C_j \right)^4$$

3. Compute the  $N, V, W, Z$  then  $A'_0, C'_0$

$$N = A_2 - (2^{n_1} \prod_{i=0}^{n_1-1} C_i)^2 x$$

$$V = A_2 + (2^{n_1} \prod_{i=0}^{n_1-1} C_i)^2 x$$

$$W = C_2 - (2^{n_1} \prod_{i=0}^{n_1-1} C_i)^3 y$$

$$Z = (2^{k_1} \prod_{i=0}^{k_1-1} C_i) N$$

$$A'_0 = W^2 - VN^2$$

$$C'_0 = -Z^3 y - W(A'_0 - Z^2 x)$$

4. if  $(n_2 > 0)$  Compute  $B'_0$ ,  $B'_0 = 3A'_0{}^2 + aZ^4$

For  $i$  from 1 to  $n_2$  Compute  $A'_i, C'_i$ , for  $i$  from 1 to  $n_2 - 1$  Compute  $B'_i$

$$A'_i = B'_{i-1}{}^2 - 8A'_i C_i{}^2$$

$$C'_i = -8C'_{i-1}{}^4 - B'_{i-1}(A'_i - 4A'_i C_i{}^2)$$

$$B'_i = 3A'_{i-1}{}^2 + 16^i aZ^4 \left( \prod_{j=0}^{i-1} C'_j \right)^4$$

Compute  $Z$ ,  $Z = Z(2^{n_2} \prod_{i=0}^{n_2-1} C'_i)$

5. Compute  $x'_{2^{k_2}}, y'_{2^{k_2}}$

$$x'_{2^{n_2}} = \frac{A'_{n_2}}{Z^2}$$

$$y'_{2^{n_2}} = \frac{C'_{n_2}}{Z^3}$$

Theorem 3.1 describes the computational complexity of this formula.

**Theorem 3.1** *In terms of affine coordinates, there exists an algorithm that computes  $2^{n_2}(2^{n_1}P + Q)$  at most  $[4(n+2) + 2]M$ ,  $[4(n+1) + 2]S$ , and  $I$  in  $\mathbf{F}_p$  for any point  $P, Q \in E(\mathbf{F}_p)$  where  $M, S$  and  $I$  denote multiplication, squaring and inversion respectively, and  $n = n_1 + n_2$ .*

**Proof** The complexity of step 1 and step 2 the same as in [12, Algorithm1] involve  $(2M + 3S)n_1 + (M+S)(n_1 - 1) + S$

In step 3, we first compute  $\prod_{i=0}^{n_1-1} C_i$  which takes  $n_1 - 1$  multiplication. Secondly, we perform one squaring to compute  $(2^{n_1} \prod_{i=0}^{n_1-1} C_i)^2$ . Next, we perform one multiplication to compute  $(2^{n_1} \prod_{i=0}^{n_1-1} C_i)^2 x$ . Then we obtain  $N$ , and  $V$ . Next, we perform two multiplications, one multiplication to compute  $(2^{n_1} \prod_{i=0}^{n_1-1} C_i)^2 y$  and other to

compute  $(2^{n_1} \prod_{i=0}^{n_1-1} C_i)(2^{n_1} \prod_{i=0}^{n_1-1} C_i)^2 y = (2^{n_1} \prod_{i=0}^{n_1-1} C_i)^3 y$ . Then we obtain  $W$ . Third we perform two squaring to compute  $W^2, N^2$ , and one multiplication to compute  $VN^2$ . Then we obtain  $A'_0$ . Forth, we perform one multiplication to compute  $(2^{n_1} \prod_{i=0}^{n_1-1} C_i)N$ . Then we obtain  $Z$ . Next we perform two squaring to compute  $Z^2, Z^4$ , and one multiplication to compute  $Z^3$ . Next we perform two multiplications to compute  $Z^2x, z^3y$ , Finally we perform one multiplication to compute  $W(A'_0 - Z^2x)$ . Then we obtain  $C'_0$ . The complexity of step 3 involves  $(n_1 - 1)M + 9M + 5S$ .

In step 1 we perform one squaring to compute  $A'_0{}^2$ . Next we perform one multiplication to compute  $aZ^4$ , where  $Z^4$  is computed in step 3. Then we obtain  $B'_0$ . The complexity of step 4.1 involve  $M + S$  and the complexity of step 2 involves  $(2M + 3S)n_2 + (M+S)(n_2 - 1)$  as step 2.

In step 3 we compute  $\prod_{i=0}^{n_2-1} C'_i$  which takes  $n_2 - 1$  multiplications. Secondly, we perform one multiplication to compute  $Z(2^{n_2} \prod_{i=0}^{n_2-1} C'_i)$ . Then we obtain new value for  $Z$ . the complexity of sub-step 3 involves  $n_2 M$ . Hence, the complexity of step 4 involves  $4n_2 M + 4n_2 S$ .

In step 5, we perform one inversion to compute  $Z^{-1}$  and the result is set to  $T$ . Next, we perform one squaring to compute  $T^2$ . Next, we perform one multiplication to compute  $A'_{n_2} T^2$ . Then we obtain  $x'_{2^{n_2}}$ . Finally we perform two multiplications to compute  $C'_{n_2} T^2 T$ . Then we obtain  $y'_{2^{n_2}}$ . The complexity of step 5 involves  $3M + S + I$ . So the complexity of above computations involve  $[4(n+2) + 2] M, [4(n+1) + 2] S$ , where  $n = n_1 + n_2$ . ■

### 3.2 The Break-Even Point

For application in practice it is highly relevant to compare the complexity of our newly derived formulae for direct computing of  $n$  doublings separated with one addition and individual  $d$  doublings. The performance of the new method depends on the cost factor of one inversion relatively to the cost of one multiplication. For this purpose, we introduce, as [5], the notation of a "break even point." It is possible to express the time that it takes to perform one inversion in terms of the equivalent number of multiplication needed per inversion. Table 3.1 shows the number of squarings  $S$ , multiplications  $M$ , and inversions  $I$  in  $\mathbf{F}_p$ .



Table 3.1 Complexity comparison: Individual doublings and one addition vs. direct computation of several doublings with one addition.

Calculation with n	Method	Complexity			Break-Even Point
		<i>S</i>	<i>M</i>	<i>I</i>	
4	DECDBL(4)	22	26	1	$6.6 M < I$
	4 doublings + 1 addition	10	9	5	
5	DECDBL(5)	26	30	1	$6 M < I$
	5 doublings + 1 addition	12	11	6	
<i>w</i>	DECDBL( <i>w</i> )	$4w+6$	$4w+10$	1	$\frac{(3.6 w + 12)}{w} M$
	<i>w</i> doublings + 1 addition	$2w+1$	$2w+2$	$w+1$	

In general let  $n = n_1 + n_2$ , let us denote the direct computing of  $2^{n_2} (2^{n_1} P + Q)$  by symbol DECDBL(*n*). Then our formulae can outperform the regular double and add algorithm if the following relation to hold:

$$\text{Cost( separate } n \text{ ECDBL + ECADD)} > \text{Cost( DECDBL}(n))$$

Ignoring squarings and additions and expressing the Cost function in terms of multiplications and inversions, we have:

$$(2n M + 2n S + n I + 2M + S + I) > (4(n+2)M + 4(n+1)S + 2M + 2S + I)$$

We define  $r = I/M$  (the ratio of speed between a multiplication and inversion), and assume that one squaring has complexity  $S = 0.8 M$  [12]. We also assume that the cost of field addition and multiplication by small constants can be ignored. One can rewrite the above expressions as:

$$n r M > (2nM + 8M + 1.6n M + 4M)$$

Solving for *r* in terms of *M* one obtains:

$$r > \frac{(3.6 n + 12)}{n} M$$

As we can see from Table 3.1, if a field inversion has complexity  $I > 7.6 M$ , direct computation of 3 doublings with one addition may be more efficient than 3 separate doubling and one adding.

### 3.2 Exponentiation with Direct Computation of $2^{n_2} (2^{n_1} P + Q)$

By using our previous formulae for direct computation of  $2^{n_2} (2^{n_1} P + Q)$ , where  $n_1 \geq 1$ , and  $P, Q \in E(\mathbb{F}_p)$ , we can improve algorithm B.1 [11] for elliptic curve exponentiation with *w*MOF by change the step 3.2 of algorithm B.1 [11] with a new step that compute  $2^{n_2} (2^{n_1} P + Q)$  directly as in the following algorithm.

---

#### **Algorithm 3.2 Exponentiation with *w*MOF Using Direct Computation of $2^{n_2} (2^{n_1} P + Q)$**

---

INPUT a non-zero *t*-bit binary string *k*,  $P \in E(\mathbb{F}_p)$ , and the multiple of the point *P*,  $\gamma_{0...tw}$  and  $\xi_{0...tw}$ , the precomputed table look-up.

OUTPUT exponentiation *kP*.

1.  $i \leftarrow t$
2.  $Q \leftarrow O$
3. While  $i \geq 1$  do the following

- 3.1. if  $(k_i \text{ XOR } k_{i-1}) = 0$ , then do the following
    - 3.1.1.  $Q \leftarrow \text{ECDBL}(Q)$
    - 3.1.2.  $i \leftarrow i - 1$
  - 3.2. else do the following
    - 3.2.1.  $\text{index} \leftarrow ((k \gg (i - w)) \& (2^{w+1} - 1)) - 2^{w-1}$
    - 3.2.2. if  $(i < w)$   $Q \leftarrow 2^{i-(w-\xi_{\text{index}})+1} (2^{w-\xi_{\text{index}}} Q + \gamma_{\text{index}} P)$
    - 3.2.3 else if  $(i \geq w)$   $Q \leftarrow 2^{\xi_{\text{index}}} (2^{w-\xi_{\text{index}}} Q + \gamma_{\text{index}} P)$
    - 3.2.4.  $i \leftarrow i - w$
  4. If  $i = 0$  do the following
    - 4.1.  $Q \leftarrow \text{ECDBL}(Q)$
    - 4.2. If  $k_0 = 1$  then  $Q \leftarrow \text{ECADD}(Q, -P)$
  5. return  $Q$
- 

In algorithm 3.2, for each window width  $w$  of  $w\text{MOF}$ , Step 3.1 performs direct computation of  $2^{i-(w-\xi_{\text{index}})+1} (2^{w-\xi_{\text{index}}} Q + \gamma_{\text{index}} P)$  if  $(i < w)$  otherwise Step 3.2 performs direct computations of  $2^{\xi_{\text{index}}} (2^{w-\xi_{\text{index}}} Q + \gamma_{\text{index}} P)$  if  $(i \geq w)$ , where  $\xi_{\text{index}} = 0, 1, \dots, w-1$ ,  $\gamma_{\text{index}} P = \{\pm 1, \pm 3, \dots, \pm(2^{w-1} - 1)\}$ .

### 3.2 Complexity Analysis of the $w\text{MOF}$ Method

In this subsection, we perform an analysis of  $w\text{MOF}$  method when it used in conjunction with the  $2^{n_2} (2^{n_1} P + Q)$  formula. In addition, we compare the complexity of  $w\text{MOF}$  method, with and without formula. Moreover we derive an expression that predicts the theoretical improvement of the  $w\text{MOF}$  method by using the formulae, in terms of the ratio between inversion and multiplication times.

Theorem 3.2 describes the complexity of algorithm B.1 [11] for computing exponentiation with  $w\text{MOF}$ .

**Theorem 3.2** *In terms of affine coordinate, let  $P \in E(\mathbb{F}_p)$ ,  $t$ -digits exponent in  $w\text{MOF}$ , then the complexity of algorithm B.1 [11] for computing  $kP$  requires on average  $\frac{(2w+4)t}{w+1}M + \frac{(2w+3)t}{w+1}S + \frac{(w+2)t}{w+1}I$ , where  $M$ ,  $S$  and  $I$  denote multiplication, squaring and inversion respectively.*

**Proof** We noticed that algorithm B.1 [11] performs an ECADD operation each time the current digit  $\delta$  is non-zero, recall from theorem 4 [11] that the average non-zero density of  $w\text{MOF}$  is asymptotically  $\frac{1}{w+1}$  also, one ECDBL operation is performed in each iteration (where  $i \geq 0$ ) to double the intermediate result. Then on average, algorithm B.1 [11] for computing exponentiation with  $w\text{MOF}$  requires

$$t \text{ ECDBL} + \frac{t}{w+1} \text{ ECADD}$$

Recall that the computational costs for doubling and additions operations in affine coordinate. Then we can rewrite previous expression as:

$$(2M + 2S + I)t + \frac{t}{w+1}(2M + S + I)$$

We can rewrite previous expression in terms of M, S, and I as:

$$\frac{(2w+4)t}{w+1}M + \frac{(2w+3)t}{w+1}S + \frac{(w+2)t}{w+1}I \quad \blacksquare$$

Now Theorem 3.3 describes the complexity of algorithm 3.1 for computing exponentiation with wMOF by using  $2^{n_2} (2^{n_1} P + Q)$ .

**Theorem 3.3** *In terms of affine coordinate, let  $P \in E(\mathbb{F}_p)$ , and  $t$ -digits exponent in wMOF, then the complexity of algorithm 3.1 for computing  $kP$  requires on average  $\frac{4(w+3)t}{w+1}M + \frac{4(w+2)t}{w+1}S + \frac{2t}{w+1}I$ , where  $M$ ,  $S$  and  $I$  denote multiplication, squaring and inversion respectively.*

**Proof** Recall from theorem 4 [11] that for  $t$ -digits exponent  $k$  in its wMOF, if  $t \rightarrow \infty$  the average non-zero density of wMOF is asymptotically  $\frac{1}{w+1}$  and wMOF of  $k$  is infinity.

Long sequence constituted from two types of blocks:

1.  $b_1 = (0)$ , length of this block is 1;
2.  $b_2 = (0^i * 0^{w-i-1})$ , length of this block is  $w$  and  $0 \leq i \leq w - 1$ ;

Then the number of block  $b_2$  equals  $\frac{1}{w+1}$  because every block  $b_2$  has a non-zero bit, and the number of block  $b_1$  equals amount of 0s in wMOF – the amount of 0s in  $b_2$  which equals

$$\frac{w}{w+1}t - (w-1)\left(\frac{1}{w+1}\right)t = \frac{t}{w+1}$$

Now, step 3.1 of algorithm 3.1 performs  $\frac{1}{w+1}t$  blocks  $b_1$  and step 3.2 performs

$\frac{1}{w+1}t$  block  $b_2$  then algorithm 3.1 for computing  $kP$  requires on average

$$\frac{t}{w+1} \text{ECDBL} + \frac{t}{w+1} \text{DECDBL}(w)$$

Recall the computational costs for doublings and additions operations in affine coordinate. Then we can rewrite previous expression as:

$$\frac{t}{w+1}(2M+2S+I + 4(w+2)M + 4(w+1)S + 2M + 2S + I)$$

We can rewrite previous expression in terms of M, S, and I as:

$$\frac{4(w+3)t}{w+1}M + \frac{4(w+2)t}{w+1}S + \frac{2t}{w+1}I \quad \blacksquare$$

### Relative Improvement

Let us denote the times it would take to perform exponentiation by using algorithms B.1 [11], and 3.1 by symbols  $T_{\text{Regular method}}$ ,  $T_{\text{Formula method}}$  respectively. According to theorems B.1 [11], and 3.1, we can derive expressions for the time it would take to perform a whole exponentiation with  $w$ MOF as:

$$T_{\text{Regular method}} = \frac{(2w+4)t}{w+1}M + \frac{(2w+3)t}{w+1}S + \frac{(w+2)t}{w+1}I \quad (3.21)$$

$$T_{\text{Formula method}} = \frac{4(w+3)t}{w+1}M + \frac{4(w+2)t}{w+1}S + \frac{2t}{w+1}I \quad (3.22)$$

From equations 3.21, and 3.22, one can readily derive the relative improvement by defining  $r = I/M$  (the ratio of speed between a multiplication and inversion) as:

$$\text{Relative Improvement} = \frac{T_{\text{Regular method}} - T_{\text{Formula method}}}{T_{\text{Regular method}}} \quad (3.23)$$

By using (3.21) and (3.22)

$$\text{Relative Improvement} = \frac{wI - [(2w+8)M + (2w+5)S]}{(w+2)I + [(2w+4)M + (2w+3)S]} \quad (3.24)$$

In our implementation  $S \approx M$  and  $r = 12.6$ , let  $w = 4$ , then

$$\text{Relative Improvement is} = \frac{4(r) - 29}{6(r) + 23} \quad (3.25)$$

$$\text{Relative Improvement is} = \frac{4(12.6) - 29}{6(12.6) + 23} \cdot 100 = 21.7\% \quad (3.26)$$

## 4. Implementation and Results

In this section, we implement our methods and others, which have been given in previous sections to show the actual performance of exponentiation. Implementation of an ECC system have several choices, these include selection of elliptic curve domain parameters, platforms [2].

### 4.1 Elliptic Curves domain parameters and Platforms

Generating the domain parameters for elliptic curve is vary time consuming. It consists of a suitably chosen elliptic curve  $E$  defined over a prime finite field  $\mathbf{F}_p$ , and a base point  $G \in E(\mathbf{F}_p)$ . Therefore we select NIST-recommended elliptic curves domain parameters in [10]. We implement 4 elliptic curves over prime fields  $\mathbf{F}_p$ , the prime modulo  $p$  are of a special type (generalized Mersenne numbers) with  $\log_2 p = 160, 192, 224, 256$ . We call these curves as P160, P192, P224, or 256 respectively.

The ECC is implemented on a Pentium 4 personal computer (PC) with 2.0 GHz processor and 512 MB of RAM. Programs were written in Java language for multi-precision integer operations, and are ran under Windows XP.

We used jBorZoi Library [1] in this implementation. jBorZoi is a Java Elliptic Curve Cryptography which implements cryptographic algorithms using elliptic curves defined over binary finite fields. We extended jBorZoi Library to implement cryptographic algorithms using elliptic curves defined over prime finite fields  $\mathbf{F}_p$ .

### 4.2 Timings analysis of wMOF Exponentiation Method

We performed timing measurements on the individual  $k$  doublings and one addition operations and the corresponding formulae for direct computation of one addition adjoint with  $k$  doublings. In addition, we developed timing estimates based on the approximately ratio of speed between a multiplication and inversion  $I / M$  in prime field  $F_p$  as presented in Table 4.1.

Table 4.1 The ratio of speed between a multiplication and inversion in prime field  $F_p$

Curves	Average Timing ( $\mu$ sec) for M	Average Timing ( $\mu$ sec) for S	Average Timing ( $\mu$ sec) for I	$r = I / M$
P160	7.0	6.9	88.0	12.6
P192	8.7	8.6	108.8	12.5
P224	10	9.8	123.1	12.3
P256	11.9	11.8	145.2	12.2

#### 4.2.1 Optimal Window Size

To show the actual improvement of wMOF method with our new formula, we must find out the most efficiency proper window size, where the length of input binary form is 160-bits, 192-bits, 224-bits, or 256-bits. Figures (4.1- 4.4) illustrate the relation among the window size  $w$ , the speed of the evaluation and pre-computed processes. We can noticed from these Figures that when the window size increases, time of the evaluation will decrease, while time of the precomputation will increase, and the optimal  $w$  is 4 when the input is 160-bits. and the optimal  $w$  is 5 when the inputs is 192, 224 or 256-bits. So all the tests in this thesis will be processed for  $w = 4$  in 160-bits input and  $w = 5$  for 192, 224, or 256-bits.




Figure 4.1 Pre-compute and evaluation with 160-bits input




Figure 4.2 Pre-compute and evaluation with 192-bits input




Figure 4.3 Pre-compute and evaluation with 224-bits input




Figure 4.4 Pre-compute and evaluation with 256-bits input

### 4.2.2 The performance of improved wMOF method

Using Table 4.1, we can readily predict that the timings for performing an exponentiation with and without the formulae presented in Algorithm 3.1. In addition, using the complexity shown in equations (3.21, 3.22) and the timings shown in Table 4.1 we can make estimates as to how long an exponentiation with wMOF will take using both doublings with formulae and individual doublings.

Table 4.2 Average time comparison required to perform an exponentiation without pre-computations stage of a random point in msec (Pentium IV 2.0 GHz)

Curves	Method	Predicted Timing	Measured Timing	% Improvement	
				Predicted	Measured
P 160	wMOF with formula (w = 4)	17.4	18.3	21.62	21.8
	wMOF (w = 4)	22.2	23.4		
P 192	wMOF with formula (w = 5)	23.8	24.3	25.62	25.7
	wMOF (w = 5)	32	32.6		
P 224	wMOF with formula (w = 5)	31.7	33.9	24.52	24.6
	wMOF (w = 5)	42	45		
P 256	wMOF with formula (w = 5)	43.8	47.4	23.5	23.3
	wMOF (w = 5)	57.3	61.8		

### Conclusion

In this paper, we construct efficient algorithm for exponentiation on elliptic curve defined over  $F_p$  in terms of affine coordinates. The algorithm computes  $2^{d_2}(2^{d_1}P+Q)$  directly from random points P and Q on an elliptic curve, without computing the intermediate points. Moreover, we apply the algorithm to exponentiation on elliptic curve with wMOF and analyze their computational complexity. A comparison was made based on notation of a "break even point." which is the cost factor of one inversion relatively to the cost of one multiplication.

This algorithm can speed the wMOF exponentiation of elliptic curve of size 160-bit about (21.7%) as a result of its implementation with respect to affine coordinates.

## References

- [1] borzoi 1.02 - an open source Elliptic Curve Cryptography Library by Dragongate Technologies Ltd., April 2004. <http://www.dragongate-technologies.com>.
- [2] M. Brown, D. Hankerson, J. Lopez, A. Menezes, Software Implementation of the NIST Elliptic Curves Over Prime Fields, Topics in Cryptology - CT-RSA 2001, LNCS 2020 (2001) 250-265.
- [3] H. Cohen, A. Miyaji, T. Ono, Efficient Elliptic Curve Exponentiation Using Mixed Coordinates, Advances in Cryptology – ASIACRYPT '98, LNCS 1514, Springer (1998) 51-65.
- [4] W. Diffie, M. Hellman, New directions in cryptography, IEEE Transactions on Information Theory, IT-22(6) (1976) 644-654.
- [5] J. Guajardo, C. Paar, Efficient Algorithms for Elliptic Curves Cryptosystem, Advances in Cryptography-CRYPTO'97, LNCS, 1294(1997), Springer-Verlage 342-356.
- [6] K. Koyama, Y. Tsuruoka, Speeding Up Elliptic Curve Cryptosystems using a Signed Binary Windows Method, Advances in Cryptology-CRYPTO'92, LNCS740 (1992) 345-357.
- [7] V.S. Miller, Use of Elliptic Curves in Cryptography, Advances in Cryptology - CRYPTO'85, LNCS 218, Springer (1986) 417-426.
- [8] A. Miyaji, T. Ono, H. Cohen, Efficient Elliptic Curve Exponentiation, Information and Communication Security - ICICS 1997, LNCS 1334, Springer (1997) 282-291.
- [9] B. Moller, Improved Techiques for Fast Exponentiation Information Security and Cryptology - ICISC 2002, LNCS 2587, Springer (2003) 298-312.
- [10] National Institute of Standard and Technology, Digital Signature Standard, FIPS Publication 186-2, February 2000.
- [11] K. Okeya, K. Schmidt-Samoa, C. Spahn, T. Takagi, Signed Binary Representations Revisited, Advances in Cryptology – CRYPTO 2004, LNCS 3152, Springer (2004) 123-139.
- [12] Y. Sakai, K. Sakurai, Efficient Scalar Multiplications on Elliptic Curves with Direct Computations of Several Doublings. IEICE Trans. Fundamentals, E84-A(1) (2001) 120-129.